§51.1565

TABLE IV—FLATTENED OR DEPRESSED AREAS—PRESSURE BRUISES MAXIMUM AREA ALLOWED

| Diameter | Weight | No. 1 (aggregate area) | No. 2 (aggregate area) |
|--|----------------------------|---------------------------|--|
| Potato is: Less than 2 in 2 to 2½ in More than 2½ to 3 in More than 3 to 3½ in More than 3½ to 4 in More than 4½ to 4½ in More than 4½ to 5 in More than 5 in | Potato is: Less than 4 oz | Not more than: 1/2 in | Not more than: 1 in 1½ in 1½ in 1¾ in 1⅙ in 2 in 2¼ in 2¾ in 3¼ in |

TABLE V—DEPTH ALLOWED FOR GROWTH CRACKS

| Diameter | Weight | No. 1 (depth) | No. 2 (depth) |
|---|--------------|-----------------------|--|
| Potato is: Less than 2 in 2 to 2½ in More than 2½ to 3 in More than 3 in | 4 oz to 6 oz | Not more than: 1/8 in | Not more than: 1/4 in 3/6 in 1/2 in 5/6 in |

TABLE VI—SUNKEN DISCOLORED AREAS MAXIMUM AREA ALLOWED

| Diameter | Weight | No. 1 (aggregate area) | No. 2 (aggregate area) |
|--|----------------------------|---------------------------|--|
| Potato is: Less than 2 in 2 to 2½ in More than 2½ to 3 in More than 3 to 3½ in More than 3½ to 4 in More than 4½ to 4½ in More than 4½ to 5 in More than 5 in | Potato is: Less than 4 oz | Not more than: 3/6 in | Not more than: 3/4 in 1 in 1 1/4 in 11/2 in 13/4 in 2 in 21/4 in 21/2 in |

[35 FR 18258, Dec. 1, 1970, as amended at 73 FR 15055, Mar. 21, 2008; 73 FR 70586, Nov. 21, 2008]

§51.1565 Internal defects.

Internal defects are defects which cannot be detected without cutting the potato. Some internal defects are listed in Table IV.

TABLE VII—INTERNAL DEFECTS

| Defects | Damage maximum allowed | Serious damage maximum allowed | | |
|---|--|-----------------------------------|--|--|
| Occurring of | Occurring outside of or not entirely confined to the vascular ring | | | |
| Ingrown Sprouts, Internal Discoloration, Vascular Browning, Fusarium Wilt, Net Necrosis, Other Necrosis, Stem End Browning. | 5 percent waste | 10 percent waste. | | |

TABLE VII—INTERNAL DEFECTS—Continued

| Defects | Damage maximum allowed | Serious damage maximum allowed |
|---|---|--|
| Internal Black Spot. | When the spot(s) are darker than the official color chip (POT—CC—2) after removing 5 percent of the total weight of the potato. | When the spot(s) are darker than the official color chip (POT-CC-2) after removing 10 percent of the total weight of the potato. |
| Occurri | ng entirely within the | vascular ring |
| Hollow Heart or Hollow Heart with Discol- oration. | Area affected not to exceed that of a circle ½ inch in diameter in a potato 2½—inches in diameter or 6 ounces in weight. 1 | Area affected not to exceed that of a circle ¾ inch in diameter in a potato 2½-inches in diameter or 6 ounces in weight.¹ |

TABLE VII—INTERNAL DEFECTS—Continued

| Defects | Damage maximum allowed | Serious damage maximum allowed |
|---|---|---|
| Light Brown Discoloration (Brown Center). | Area affected not to exceed that of a circle ½ inch in di- ameter in a potato 2½ inches in di- ameter or 6 ounces in weight.¹ | Area affected not to exceed that of a circle ¾ inch in di- ameter in a potato 2½ inches in di- ameter or 6 ounces in weight.¹ |

TABLE VII—INTERNAL DEFECTS—Continued

| Defects | Damage maximum allowed | Serious damage maximum allowed | |
|---|--|--|--|
| Occurring entirely within the vascular ring | | | |
| Internal Brown Spot and Similar Dis- coloration (Heat Necro- sis). | Not more than the equivalent of 3 scattered spots 1/6 inch in diameter in a potato 21/2—inches in diameter or 6 ounces in weight.1 | Not more than the equivalent of 6 scattered spots ½ inch in diameter in a potato 2½-inches in diameter or 6 ounces in weight.¹ | |

¹Note: Correspondingly lesser or greater areas in smaller or larger potatoes.

[35 FR 18258, Dec. 1, 1970. Redesignated at 42 FR 32514, June 27, 1977, and further redesignated at 46 FR 63203, Dec. 31, 1981; amended at 56 FR 7553, Feb. 25, 1991; 56 FR 10302, Mar. 11, 1991; 73 FR 15058, Mar. 21, 2008; 73 FR 70586, Nov. 21, 2008]

METRIC CONVERSION TABLE

§51.1566 Metric conversion table.

| Inches | Millimeters (mm) | |
|-------------|---------------------|--|
| 1/8 equals | 3.2 | |
| 1/4 equals | 6.4 | |
| 1/2 equals | 12.7 | |
| 3/4 equals | 19.1 | |
| 1 equals | 25.4 | |
| 11/2 equals | 38.1 | |
| 2 equals | 50.8 | |
| 21/2 equals | 63.5 | |
| 3 equals | 76.2 | |
| 31/2 equals | 88.9 | |
| 4 equals | 101.6 | |
| 41/2 equals | 114.3 | |

| Ounces | Grams |
|-----------|--------|
| 1 equals | 28.35 |
| 4 equals | 113.40 |
| 5 equals | 141.75 |
| 6 equals | 170.10 |
| 7 equals | 198.45 |
| 8 equals | 226.80 |
| 9 equals | 255.15 |
| 10 equals | 283.50 |
| 12 equals | 340.20 |
| 14 equals | 396.90 |
| 16 equals | 453.60 |
| 18 equals | 510.30 |
| 19 equals | 538.60 |
| 20 equals | 567.00 |

Subpart—United States Consumer Standards for Potatoes

SOURCE: 18 FR 7132, Nov. 11, 1953, unless otherwise noted. Redesignated at 42 FR 32514, June 27, 1977, and further redesignated at 46 FR 63203, Dec. 31, 1981.

GRADES

§51.1575 U.S. Grade A Small; U.S. Grade A Medium; U.S. Grade A Medium to Large; U.S. Grade A Large.

Potatoes of each of these grades shall be of one variety or similar varietal characteristics which are fairly well shaped, fairly clean, free from freezing injury, blackheart, late blight, and soft rot or wet breakdown, and from damage caused by sunburn, second growth, growth cracks, air cracks, hollowheart, internal discoloration, cuts, shriveling, sprouting, scab, dry rot, rhizoctonia, other diseases, wireworm, other insects, or mechanical or other means. Potatoes of these grades shall also be mature: *Provided*, That potatoes which are not mature and the outer skin loosens or "feathers" readily under the usual handling practices need not meet this requirement if they are firm and are further designated as "Early" in connection with the grade, as for example "U.S. Grade A Medium-Early." Potatoes on the shown face shall be reasonably representative in size and quality of the contents of the containers. (See § 51.5177.)

(a) *Tolerances.* Incident to proper grading and handling, except for the tolerances for size, not more than a total of 5 percent, by weight, of the potatoes in any lot may fail to meet the requirements of the grade, including not more than 1 percent for potatoes affected by soft rot or wet breakdown. (See §51.1579.)